

# ABANTEL™



## BROAD SPECTRUM WORMER FOR DOGS

### COST EFFECTIVE BROAD SPECTRUM WORMER FOR DOGS

#### ACTIVE CONSTITUENTS:

Each tablet contains: 545mg OXANTEL EMBONATE,  
140mg PYRANTEL EMBONATE, and 50mg PRAZIQUANTEL

For treatment and control of roundworms (*Toxocara* spp., *Toxascaris leonina*), hookworms (*Uncinaria stenocephala*, *Ancylostoma* spp), whipworms (*Trichuris vulpis*) and all tapeworms (*Taenia* spp, *Dipylidium caninum*, *Spirometra erinacei* and *Echinococcus granulosus*), in dogs.



	Popantel®	Abantel™
Tablet Dose	10kg dog	10kg dog
Active Ingredient		
Praziquantel	50mg	50mg
Pyrantel Embonate	143mg	140mg
Oxantel Embonate	542mg	545mg
Controls Roundworms, Whipworms, Hookworms, & Tapeworms	YES	YES

Popantel® is registered trademark Jurox Pty Ltd, 85 Gardiner St, RUTHERFORD, NSW, 2320, AUSTRALIA

Abantel™ is a trademark of Abbey Animal Health Pty Ltd

# ABANTEL™

## BROAD SPECTRUM WORMER FOR DOGS

**Active Constituent:** Each tablet contains: 545mg OXANTEL EMBONATE, 140mg PYRANTEL EMBONATE, and 50mg PRAZIQUANTEL

For treatment and control of roundworms (*Toxocara* spp., *Toxascaris leonina*), hookworms (*Uncinaria stenocephala*, *Ancylostoma* spp), whipworms (*Trichuris vulpis*) and all tapeworms (*Taenia* spp, *Dipylidium caninum*, *Spirometra erinacei* and *Echinococcus granulosus*), in dogs.

This product does not control heartworm.

If worm problems persist, consult a veterinarian.

### DIRECTIONS FOR USE

#### DOSAGE AND ADMINISTRATION

For oral use only. Can be administered directly into the mouth or crushed and mixed with the feed.

1 tablet per 10kg bodyweight. Dogs up to 5kg should be given half (½) a tablet.

#### Dosage table:

Bodyweight	Dose
up to 5kg	½ tablet
10kg	1 tablet
20kg	2 tablets
30kg	3 tablets
40kg	4 tablets
50kg	5 tablets

Specific treatment for *Spirometra erinacei* is by administering ABANTEL™ BROAD SPECTRUM WORMER FOR DOGS four (4) times the above dose rate.



### GENERAL DIRECTIONS

**ROUNDWORMS & HOOKWORMS:** Treat at 2, 4, 8 and 12 weeks - thereafter every 3 months.

**WHIPWORMS:** Treat every 6-8 weeks after 3 months of age.

**HYDATID TAPEWORM:** Dog should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks.

**OTHER TAPEWORMS:** Treat every 3 months.

**PREGNANT BITCHES:** Treat at mating, before whelping and then every 3 months.

For *Dipylidium* treatment adequate flea control should be maintained all year around.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre: Phone Australia 131 126

### DISPOSAL

Dispose of empty container by wrapping with paper and placing in garbage.

### STORAGE

Store below 25°C (Air Conditioning) in a dry place.

APVMA Approval No: 69893/111735