

When your pigs need treatment
for inflammation, pain
or fever



Water them well

Tap into Australia's only NSAID medicated water solution that
delivers effective relief from inflammation, pain and fever.



Why use ANIBEX™ SOLUBLE POWDER FOR PIGS?

When a herd of pigs become impacted with viral or bacterial infections causing inflammation, pain or fever, they need quick access to an effective NSAID medication.

A broad based method of delivering ANIBEX™ via the water supply provides timely relief to the group. It is more effective and less stressful for both the animals and care giver to administer medications via the water supply.¹



What are the benefits of ANIBEX™ SOLUBLE POWDER FOR PIGS?



Contains sodium salicylate, which dissolves readily in drinking water,² providing NSAID medicated water for the herd.



Rapidly absorbed into the bloodstream.²



Effective in the treatment of inflammation in pigs, including pain, fever and inflammation caused by viral or bacterial infections.



Can be used concomitantly with antibiotics when required.



Sodium salicylate has been tried and tested in Europe for over 30 years.

Solubility

Sodium salicylate is highly soluble in water² which means concentrated stock solution can be made within just a few seconds.

Sodium salicylate remained soluble for a full 24 hour period under both laboratory and nursery testing conditions.²

It has been proven that there was no interaction of sodium salicylate with any other substances in drinking water that could affect the bioavailability and absorption of sodium salicylate in pigs.²

Dosage and Administration

ANIBEX™ is administered orally in drinking water.

The recommended dose rate is 35 mg/kg body weight per day once daily for a maximum of 5 days, orally in drinking water.

Mixing Directions

Individual tank or Bulk tank use: Water consumption should be calculated on daily basis or user should seek information from veterinarian/ industry recommendations about average water intake per pigs' age or weight range. Water consumption in pigs can vary depending on the pigs' age and size, temperature difference and seasons.

The following formula can be used to calculate the amount of ANIBEX™ in drinking water:

Individual or Bulk tank formula

$$\frac{35 \text{ mg ANIBEX}^{\text{TM}}/\text{kg bodyweight/day} \times \text{Mean body weight (kg) of pigs to be treated}}{\text{Mean daily water (L) per animal}} = \text{___ mg ANIBEX}^{\text{TM}} \text{ per L drinking water}$$

Multiply the calculated amount of mg ANIBEX™ per L x the total L required for the individual or bulk tank use.

Proportioner: Determine the dose using the formula below. Mix the dose of ANIBEX™ and water in the medication tank. The amount of water in the medication tank should be 20 to 100L. The medicated water should be administered to the water line via a proportioner over a period of up to 24 hours.

Proportioner formula

$$35 \text{ mg ANIBEX}^{\text{TM}}/\text{kg bodyweight/day} \times \text{Mean body weight (kg) of pigs to be treated} \times \text{Number of pigs treated} = \text{___ mg ANIBEX}^{\text{TM}}$$

ANIBEX™ SOLUBLE POWDER FOR PIGS

ACTIVE CONSTITUENT: 1000g/kg SODIUM SALICYLATE

For the symptomatic treatment of fever and inflammation in pigs.



INDICATIONS

ANIBEX™ is effective in the treatment of inflammation in pigs, and also effective in the treatment of pain during fever and inflammation caused by viral or bacterial infections. ANIBEX™ can be used in combination with concurrent antibiotic therapy in pigs.

DESCRIPTION

Sodium salicylate inhibits the biosynthesis of natural prostaglandins. By inhibiting prostaglandin synthesis, inflammation and pain perception will be reduced. Sodium salicylate has an antipyretic, anti-inflammatory and analgesic mode of action. It is absorbed quickly in the gastrointestinal tract. Sodium salicylate is metabolised in the liver and excreted via urine and milk.

DIRECTIONS FOR USE

Contraindications

This product should not be used in animals with known hypersensitivity to sodium salicylate. This product should not be used in combination with drugs known to have anticoagulant properties. This product should not be used in animals with severe liver and kidney affections. This product should not be used in case of gastrointestinal ulcerations and chronic gastrointestinal disorders. This product should not be used in case of malfunction of the hematopoietic system, coagulopathies, and haemorrhagic diathesis.

Precautions

DO NOT use in piglets of less than 4 weeks of age.

DO NOT use during pregnancy and lactation. Concurrent use with any other NSAID or steroids should be avoided. Concurrent administration of potentially nephrotoxic drugs (e.g. aminoglycosides) should be avoided. Salicylic acid is highly plasma (albumin) bound and competes with a variety of compounds for plasma protein binding sites (e.g. ketoprofen). Sodium salicylate may inhibit clotting of blood, it is recommended that elective surgery should not be performed on animals within 7 days after the end of the treatment.

Side Effects

Gastrointestinal irritation may occur, especially in animals with pre-existing gastrointestinal disease. Such irritation may clinically be manifested by production of black manure due to blood loss in the gastrointestinal tract. Inhibition of normal blood clotting may occur incidentally. If this effect occurs it will be reversible and effects will diminish within approximately 7 days.

DOSAGE AND ADMINISTRATION

ANIBEX™ soluble powder can be administered orally through the drinking water. Clean drinking water should be used. Prepare fresh solutions daily. The recommended dose rate is 35 mg/kg body weight per day once daily for a maximum of 5 days.

Mixing Directions

Individual tank or Bulk tank use: Water consumption should be calculated on daily basis or user should seek information from veterinarian/ industry recommendations about average water intake per pigs' age or weight range. Water consumption in pigs can vary depending on the pigs' age and size, temperature difference and seasons.

Following formula can be used to calculate the concentration of ANIBEX™ in drinking water:

$$\frac{35 \text{ mg ANIBEX}^{\text{TM}}/\text{kg bodyweight/day} \times \text{Mean body weight (kg) of pigs to be treated}}{\text{Mean daily water (L) per animal}} = \text{mg ANIBEX}^{\text{TM}} \text{ per L drinking water}$$

Multiply the calculated amount of mg ANIBEX™ per L x the total L required for the individual or bulk tank use.

Proportioner: Determine the dose using the formula below. Mix the dose of ANIBEX™ and water in the medication tank. The amount of water in the medication tank should be 20 to 100L.

References: 1. Australian Pork Fact Sheet; 'Water Supply to Pigs' <https://australianpork.com.au/wp-content/uploads/2013/09/Fact-sheet-water.pdf> [Date accessed 26 June 2019] 2. Data on file

The medicated water should be administered to the water line via a proportioner over a period of up to 24 hours. The proportioner and medication tank should be supervised regularly to ensure the medication is being dispensed to the water line.

Following formula can be used to calculate the amount of ANIBEX™ in drinking water treated group housing:

$$35 \text{ mg ANIBEX}^{\text{TM}}/\text{kg bodyweight/day} \times \text{Mean body weight (kg) of pigs to be treated} \times \text{Number of pigs treated} = \text{mg ANIBEX}^{\text{TM}}$$

General Directions

ANIBEX™ is readily dissolved and dispersed in water. Prepare fresh solution daily and ensure that full dose is consumed within 24 hours. During treatment, do not supply liquid which is not medicated until the medicated liquid is consumed.

It is recommended during the treatment period daily water consumption is monitored to ensure correct dosing and required daily water intake. Significant variations in water consumption should be reported to the treating veterinarian. In cases where animals do not have adequate water intake or are debilitated, alternative dose method for treatments of pain and fever should be selected.

WITHHOLDING PERIOD

MEAT: DO NOT USE less than 2 days before slaughter for human consumption.

TRADE ADVICE - ESI: DO NOT USE - less than 2 days before export. Before using this product confirm the current ESI from Abbey Laboratories on 02 8088 0720 or the APVMA website.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat. Avoid contact with eyes and skin. Avoid inhaling dust. When opening the bag and preparing product for use, wear disposable gloves, goggles and disposable dust face mask covering mouth and nose. Wash hands after use. After each day's use, wash goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126. If swallowed, do NOT induce vomiting. If in eyes wash out immediately with water.

ADDITIONAL USER SAFETY INFORMATION

People with known hypersensitivity to sodium salicylate or related drugs (for example aspirin) should avoid contact with the product.

DISPOSAL

Shake container into the medicated water. Do not dispose of undiluted chemicals on-site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 25°C (air conditioning) in a dry place. Protect from light.

PACK SIZE: 5kg

ANIBEX™ is a trademark of Abbey Laboratories Pty Ltd
APVMA Approval No: 85682/113832

ABBEY LABORATORIES PTY LTD
DISTRIBUTED BY: **ABBEY ANIMAL HEALTH PTY LTD**, 16 VOYAGER CIRCUIT
GLENDEENING NSW 2761. INFOLINE: 02 8088 0720 | www.abbeylabs.com.au