

COXI-STOP™

COCCIDIOCIDES FOR PIGLETS AND CATTLE 50

ACTIVE CONSTITUENT: 50g/L TOLTRAZURIL

At Last! An Alternative Treatment for the Control of Coccidiosis in Piglets and Young Cattle!

For the control of coccidiosis in neonatal piglets caused by *Isospora suis*, and in cattle up to 9 months of age caused by *Eimeria bovis* or *Eimeria zuernii*

- Low Volume 1mL dose for piglets
- Contains the active Toltrazuril
- Can be used in piglets from 3-6 days of age
- Can be used in calves up to 9 months of age
- Easy to use suspension



	Coxi-Stop™ 50	Baycox® Piglet & Calf
Active Ingredient	Toltrazuril 50g/L	Toltrazuril 50g/L
Suspension Formulation	YES	YES
Can be used in Piglets and Calves	YES	YES
Withholding Period: Pigs	70 days	70 days
Withholding Period: Calves	56 days	56 days
ESI: Cattle	75 days	75 days
Dose Rate: Piglets	1 mL	1mL
Dose Rate: Calves	3mLs per 10kg	3mLs per 10kg

Baycox is a registered Trademark of Bayer Intellectual Property GmbH, Alfred-Nobel-Strasse 10, 40789 Monheim am Rhein, GERMANY

Coxi-Stop™ is a trademark of Abbey Laboratories Pty Ltd.

COXI-STOP™

COCCIDIOCIDES FOR PIGLETS AND CATTLE 50

ACTIVE CONSTITUENT: 50g/L TOLTRAZURIL

FOR THE CONTROL OF COCCIDIOSIS IN NEONATAL PIGLETS CAUSED BY *ISOSPORA SUIS*, AND IN CATTLE UP TO 9 MONTHS OF AGE CAUSED BY *EIMERIA BOVIS* OR *EIMERIA ZUERNII*

DIRECTIONS FOR USE

Restraints:

DO NOT RE-TREAT calves for 75 days after the last treatment.

DO NOT TREAT piglets later than 8 days of age.

DOSAGE & ADMINISTRATION

SHAKE THE CONTAINER BEFORE USE.

The product is administered orally. Use the measuring cup to give the full dose by mouth.

DOSE RATE TABLE:-

	Dose	Prevention	Treatment
Piglets	1mL	Treat at 3 - 6 days old	Treat at onset of coccidial diarrhoea. Treat all piglets in infected litters and all infected and 'at risk' litters in the farrowing house.
Cattle	3mL per 10kg bodyweight	In the case of history of coccidiosis on farm, treat 1 week before expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves, treat at weaning time when milk feeding ceases.	Treat at onset of diagnosed clinical signs.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT (PIGS): DO NOT USE less than 70 days before slaughter for human consumption.

MEAT (CATTLE): DO NOT USE less than 56 days before slaughter for human consumption.

MILK (CATTLE): DO NOT USE in lactating or pregnant cows where milk may be used or processed for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI) PIGS: An ESI established has not been established for this product for pigs. Note- observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Abbey Animal Health Pty Ltd on 02 8088 0720 before using this product.

EXPORT SLAUGHTER INTERVAL (ESI) CATTLE: DO NOT slaughter cattle for export for 75 days after last treatment. The ESI for cattle on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 02 8088 0720 or the APVMA website: www.apvma.gov.au/residues.

SAFETY DIRECTIONS

When using this product wear rubber gloves. After each day's use wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ENVIRONMENTAL PROTECTION

Do not contaminate ponds, waterways or drains with this product or used container.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval No: 81911/104400

Coxi-Stop™ is a trademark of Abbey Laboratories Pty Ltd.