

MELOXXI™ INJECTION




Active Constituent: 20 mg/mL MELOXICAM

Prescription pain relief you can trust

A non-steroidal anti-inflammatory, analgesic, antipyretic for use in cattle, pigs and horses.


- Registered for use in cattle, pigs and horses
- Short WHP and ESI
- Registered for use IV, IM or SC



	MELOXXI™ INJECTION	Metacam®
ACTIVE CONSTITUENT	20mg/mL Meloxicam	20mg/mL Meloxicam
SPECIES	Cattle, Pigs & Horses	Cattle, Pigs, Sheep & Horses
WITHHOLDING PERIODS		
Milk	6 days (12 milkings)	6 days (12 milkings)
Meat: Cattle	8 days	8 days
Meat: Pigs	4 days	4 days
ESI		
Cattle	No ESI established	No ESI established
Pigs	4 days	4 days
ROUTE OF ADMINISTRATION		
Cattle	SC or IV	SC or IV
Pigs	IM	IM
Horses	IV	IV
STORAGE AFTER BROACHING VIAL	Discard the unused portion after 28 days of first broaching the vial.	Discard the unused portion after 28 days of first broaching vial.

MELOXXI™ is a trademark of Abbey Laboratories Pty Ltd.

Metacam® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH



MELOXXI™ INJECTION

Active Constituent:
20 mg/mL MELOXICAM

A non-steroidal anti-inflammatory, analgesic, antipyretic for use in cattle, pigs and horses.

INDICATIONS

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, antiendotoxic, anti-exudative, analgesic and antipyretic properties.

Cattle

For use in acute respiratory infection and diarrhoea in combination with appropriate antibiotic therapy to reduce clinical symptoms in calves and young cattle.

For use in acute mastitis, in combination with antibiotic therapy, as appropriate, to reduce clinical symptoms in lactating cows.

For use to assist in the control of pain following the dehorning of cattle particularly that following heat cautery dehorning of young cattle. It is recommended that the injection be administered approximately 10 minutes before dehorning and be accompanied by a cornual nerve block anaesthesia.

Pigs

For use in acute non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For use in puerperal septicaemia and toxæmia (mastitismetritis-agalactia syndrome) with appropriate antibiotic therapy to reduce clinical signs of inflammation, oppose the effects of endotoxins and hasten recovery.

Horses

For single dose rapid initiation of therapy of musculoskeletal disorders and relief of pain associated with colic.

DIRECTIONS FOR USE

Contraindications

This product is contraindicated for use in animals suffering from haemorrhagic gastrointestinal disorders, impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of individual hypersensitivity to the product.

Precautions

Pigs: Safe use in pregnant sows has not been fully evaluated but preliminary studies suggest that it has no harmful effects in the second and third trimester of pregnancy. This product can be used in lactating sows.

Horses: The safety of this product in pregnant or lactating horses and in horses under 6 weeks of age has not been established.

The product should not be used concurrently with glucocorticoids, other non-steroidal anti-inflammatory drugs or other anti-coagulant agents. Use with caution in conjunction with other highly protein-bound drugs.

Side effects

Subcutaneous as well as intravenous administration is well tolerated in cattle; only a slight transient swelling at the injection site following subcutaneous administration was observed in less than 10% of the animals treated in clinical studies.

In the case of overdosage, symptomatic treatment should be initiated.

DOSAGE AND ADMINISTRATION

Discard the unused portion after 28 days of first broaching the vial.

CATTLE: SINGLE USE ONLY by subcutaneous or intravenous injection. 0.5 mg Meloxicam/kg bodyweight (i.e. 2.5 mL/100 kg bodyweight) in combination with antibiotic therapy, as appropriate.

PIGS: For SINGLE intramuscular injection in the anterior half of neck. (Can be repeated ONCE after 24 hours if necessary). 0.4 mg Meloxicam/kg bodyweight (i.e. 2.0 mL/100 kg bodyweight) in combination with antibiotic therapy, as appropriate.

HORSES: SINGLE USE ONLY by intravenous injection. 0.6 mg Meloxicam/kg bodyweight (i.e. 3.0 mL/100 kg bodyweight).

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 8 days for cattle, 4 days for pigs and 28 days for horses before slaughter for human consumption.

MILK: Milk collected from cows within 6 days (12 milkings) of treatment MUST NOT BE USED for human consumption, processing, or fed to bobby calves.

CATTLE: Repeat treatments (more than ONCE), higher doses or injection into the muscle could result in residues above the MRLs unless the label withholding period is extended.

PIGS: Repeat doses (more than TWICE) or higher doses could result in residues above the MRLs unless the label withholding period is extended.

The prescribing veterinarian would need to advise on an extended withholding period.

TRADE ADVICE- EXPORT SLAUGHTER INTERVAL (ESI)

PIGS: DO NOT USE less than 4 days before slaughter for export.

CATTLE: This product does not have an ESI established.

Before using this product, confirm the current ESI from the supplier on 02 8088 0720 or the APVMA website.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

Additional information is listed in Safety Data Sheet.

DISPOSAL

Dispose of empty container by wrapping in paper and putting in garbage.

STORAGE

Store below 25°C (air conditioning). Not to be used after 28 days of opening.

May cause damage to organs through prolonged or repeated exposure.

Do not breathe fumes, mists, vapours or spray.

Do not get in eyes, on skin, or on clothing.

Do not eat, drink or smoke when using this product.

Use personal protective equipment as required.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF ON SKIN (or hair): Remove immediately all contaminated clothing. Rinse skin with water.

In case of fire, use carbon dioxide, dry chemical, foam, water fog. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, fine water spray can be used.

APVMA Approval No: 87070/117480

MELOXXI™ is a trademark of Abbey Laboratories Pty Ltd.