

TILMICABS™ AC

(TILMICOSIN PHOSPHATE) AQUEOUS CONCENTRATE



A KEY WEAPON IN THE FIGHT AGAINST PNEUMONIA IN PIGS

- Active Constituent – Tilmicosin Phosphate
- Aqueous Concentrate formulation - mixes readily with all types of water sources
- Indicated to treat APP and Mycoplasma
- Comes in an easy to use 5L container



Feature	Tilmicabs™ AC	Pulmotil® AC
Tilmicosin (as Phosphate)	250mg/mL	250mg/mL
Aqueous Concentrate Formulation	YES	YES
Dose Rate	80 mL per 100L water	80 mL per 100L water
Registered pack sizes	250mL, 1L, 2.5L, 5L and 10L	240mL and 960mL
Withholding Periods	Meat: 7 days	Meat: 7days
Restraints	USE ONLY IN SWINE USE ONLY ORALLY	USE ONLY IN SWINE USE ONLY ORALLY

Pulmotil is a registered trademark of Eli Lilly and Company (USA).

Tilmicabs is a trademark of Abbey Laboratories Pty Ltd.

TILMICABS™ AC

(TILMICOSIN PHOSPHATE) AQUEOUS CONCENTRATE

ACTIVE CONSTITUENTS:

250mg/mL Tilmicosin (as tilmicosin phosphate)

Contents: 250mL, 1L, 2.5L, 5L, 10L

Water medication for the treatment of pneumonia in pigs caused by *Actinobacillus pleuropneumoniae*, *Actinomyces pyogenes*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, and other organisms sensitive to tilmicosin.

DIRECTIONS FOR USE

Restraints:

USE ONLY IN SWINE

USE ONLY ORALLY

PRECAUTIONS:

The safety of tilmicosin has not been established in boars intended for breeding purposes.

Tilmicosin has a bitter taste. Exceeding the registered dose rate may result in reduced water intake. Monitor water intake carefully whilst medicating with Tilmicabs™ AC.

Contraindications:

DO NOT USE this product as an injectable as it contains disodium edetate. Avoid allowing horses or other equines access to drinking water containing tilmicosin.

General Directions:

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

DOSAGE AND ADMINISTRATION:

Prepared medicated drinking water should be used within 24 hours of mixing.

To be included in the drinking water at a dose rate of 200mg per litre (80mL Tilmicabs™ AC per 100L water) for five consecutive days, to provide a daily dose of 15-20mg tilmicosin per kg bodyweight.

WITHHOLDING PERIOD:

MEAT: DO NOT USE less than 7 days before slaughter for human consumption.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on 02 9477 5005 before using this product.

PRESCRIPTION ANIMAL REMEDY

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

SAFETY DIRECTIONS

Will irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with skin and eyes. Wash hands after use. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

ENVIRONMENTAL PROTECTION: sludge from waste treatment ponds must be disposed of in approved landfills or contaminated wastes. This product may accumulate in soils. DO NOT spread effluent containing tilmicosin on the same pastures or croplands in successive years. DO NOT allow wastes from pigs treated with tilmicosin to enter waterways. These wastes contain tilmicosin which is highly toxic to algae.

DISPOSAL [250mL and 1L]:

Dispose of empty containers by wrapping in paper and putting in garbage.

DISPOSAL [5L and 10L]:

Triple or (preferably) pressure rinse into the medicated water / dip/ drench etc. Do not dispose of undiluted chemicals on-site. Return clean container for recycling where this is an option or for disposal at a landfill authorised to accept that waste. If neither of these options are available, bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers should not be burned.

STORAGE

Store below 30°C (Room Temperature) in a safe place. Protect from direct sunlight.

Tilmicabs is a trademark of Abbey Laboratories Pty Ltd

APVMA Approval No: 69566/63277