


COXI-STOP™ 25

COCCIDIOCIDAL SOLUTION FOR POULTRY

**A NEW WEAPON IN
THE CONTROL OF
COCCIDIOSIS!**



**ACTIVE CONSTITUENT:
25g/L TOLTRAZURIL**

**FOR THE TREATMENT AND CONTROL
OF COCCIDIOSIS IN CHICKENS
CAUSED BY EIMERIA SPECIES**

| | Baycox® Poultry | COXI-STOP™ 25 |
|-------------------------------------|---|---|
| Concentration of Active Toltrazuril | 25g/L | 25g/L |
| Withholding Period: MEAT | 14 days | 14 days |
| Dose Rate | 7mg/kg body-weight/day | 7mg/kg body-weight/day |
| Controls Eimeria Species | <i>Eimeria tenella</i> , <i>Eimeria necatrix</i> , <i>Eimeria acervulina</i> and <i>Eimeria maxima</i> | <i>Eimeria tenella</i> , <i>Eimeria necatrix</i> , <i>Eimeria acervulina</i> and <i>Eimeria maxima</i> |

Baycox® is a registered trademark of Bayer Intellectual Property GmbH, Germany.
COXI-STOP™ is a trademark owned by Abbey Laboratories Pty Ltd

COXI-STOP™ 25 COCCIDIOCIDAL SOLUTION FOR POULTRY

ACTIVE CONSTITUENTS:

25g/L TOLTRAZURIL

FOR THE TREATMENT AND CONTROL OF COCCIDIOSIS IN CHICKENS CAUSED BY EIMERIA SPECIES

COXI-STOP™ 25 is added to the drinking water of chickens for the treatment and control of coccidiosis caused by Eimeria species including: *Eimeria tenella*, *Eimeria necatrix*, *Eimeria acervulina* and *Eimeria maxima*.

DIRECTIONS FOR USE

Restraint:

DO NOT treat replacement pullets more than twice.

DOSAGE AND ADMINISTRATION

COXI-STOP™ 25 is administered in drinking water at a dose rate of 7mg/kg bodyweight/day

and a final drinking water concentration of 75mg/L.

One mL of COXI-STOP™ 25 contains 25mg of the active ingredient Toltrazuril.

COXI-STOP™ 25 is diluted at the rate of 3 litres per 1000 litres of drinking water (3: 1000) and administered on 2 consecutive days with an 8 hour treatment period on both days.

Add the required amount of COXI-STOP™ 25 to the amount of water consumed in an 8 hour period and stir thoroughly. Allow for varying water consumption during hot weather.

Treat the birds for 8 hours each day for 2 days (2x8 hour treatments). All other sources of drinking water should be withheld during the 8 hour treatment period. Ensure full lighting conditions during treatment. Do not withhold feed during treatment.

GENERAL INSTRUCTIONS

DO NOT USE GALVANISED OR RUSTY IRON WATER TANKS AND RETICULATION SYSTEMS. COXI-STOP 25 mixes with all types of water. However in areas where the drinking water is acidic, the water should be tested with pH test strips and brought to a minimum pH of 8.5 using sodium carbonate (washing soda).

To avoid excessive foaming, add COXI-STOP 25 to the water after filling the supply tank.

Broiler Treatment Programme:

To use COXI-STOP™ 25 in a clinical outbreak obtain an accurate diagnosis. Begin treatment with COXI-STOP™ 25 as soon as coccidiosis is confirmed. A treatment period of 2 days is sufficient. If required, treatment may be repeated after 5 days.

Broiler Preventive Programme:

To use COXI-STOP™ 25 as a coccidiosis preventive in broiler production, give COXI-STOP™ 25 as above on days 9 and 10, 16 and 17, and 23 and 24 of life. Do not use a coccidiostat in the feed. An additional treatment may be necessary after 35 days of age but not within 14 days of slaughter.

Layer Birds: COXI-STOP™ 25 may be used in replacement pullets and replacement layer breeder birds via the drinking water for an 8 hour period each day for two consecutive days. Treatment may be repeated after 5 days if required.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

EGGS: DO NOT USE in laying hens. DO NOT USE in replacement pullets within 8 weeks of laying onset, where eggs are to be used or processed for human consumption. Eggs laid during the withholding period MUST NOT BE USED or processed for human consumption.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131126.

SAFETY DATA SHEET (SDS)

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier, call 02 80907187.

ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Triple-rinse container into the medicated water. Do not dispose of undiluted chemicals onsite. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature) in the closed, original container, as cool as possible. Protect from light.

APVMA Approval No: 80980/102029